

THIOPHANATE-METHYL	GROUP	1	FUNGICIDE
ETRIDIAZOLE	GROUP	14	FUNGICIDE

# Banrot®

## 40 WP

### Broad Spectrum Fungicide 40% Wettable Powder

To control damping-off, root and stem rot diseases of ornamental and nursery crops caused by Pythium, Phytophthora, Rhizoctonia, Fusarium and Thielaviopsis.

#### ACTIVE INGREDIENTS:

Etridiazole† ..... 15%  
Thiophanate-Methyl†† ..... 25%

OTHER INGREDIENTS: ..... 60%  
Total 100%

†CAS # 2593-15-9

††CAS # 23564-05-8

EPA Reg. No. 58185-10

EPA Est. 39578-TX-1E

Superscript used is the first letter of run code on end of container.

## KEEP OUT OF THE REACH OF CHILDREN WARNING AVISO

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

Product of U.S.A.

BANROT® is a Registered Trademark of Everris International BV.

**Net Weight: 2 lb (907 g)**  
**E957252**

### FIRST AID

**If in Eyes:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**If on Skin or Clothing:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

**If Swallowed:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**If Inhaled:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-888-241-6409 for emergency medical treatment information.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed. Product causes substantial but temporary eye injury.

**Carefully read and follow attached labeling for directions for use.**

### PRECAUTIONARY STATEMENTS

#### Hazards to Humans and Domestic Animals

**WARNING:** Causes substantial but temporary eye injury. Harmful if absorbed through skin. Do not get in eyes or on clothing. Wear goggles or face shield. Avoid contact with skin. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals. Harmful if swallowed or inhaled. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse.

Manufactured for:  
Everris NA Inc.  
PO Box 3310  
Dublin, OH 43016



## Personal Protective Equipment (PPE)

Some materials that are chemical-resistant to this product are barrier laminate, butyl rubber  $\geq 14$  mils, nitrile rubber  $\geq 14$  mils, neoprene rubber  $\geq 14$  mils, and natural rubber  $\geq 14$  mils.

Mixers, loaders, applicators and handlers must wear:

- goggles or face shield
- coveralls over long-sleeved shirt and long pants
- chemical-resistant gloves made of any water-proof material
- chemical-resistant footwear plus socks
- a minimum of a NIOSH-approved elastomeric half mask respirator with organic vapor (OV) cartridges and a combination N1, R or P filter; OR a NIOSH-approved gas mask with OV canisters; OR a NIOSH-approved powered air-purifying respirator with OV cartridges and combination HE filters

Wear a chemical-resistant apron when mixing, loading, or cleaning equipment.

## User Safety Requirements

Discard clothing or other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

### User Safety Recommendations

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## Environmental Hazards

**This product is toxic to fish.** Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate.

Etridiazole can contaminate surface water through spray drift. Under some conditions, etridiazole may have a high potential for runoff into surface water for several weeks postapplication. These conditions include poorly draining or wet soils with readily visible slopes toward adjacent surface water, frequently flooded areas, areas overlaying extremely shallow ground water, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent waters with vegetated filter strips, and areas overlaying tile drainage systems that drain surface water.

## Physical or Chemical Hazards

Do not use or store near heat or open flame.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. Do not allow this product to drift. Do not graze treated areas. Do not feed clippings to livestock. For any requirements specific to your State or Tribe, consult the Agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

**INDOOR RESTRICTIONS:** Entry (including early entry that would otherwise be permitted under the WPS) into greenhouses, potting sheds, and other indoor areas by any person – other than a correctly trained applicator who is performing a handling task permitted by the WPS and who is wearing the required handler PPE including a respirator – is PROHIBITED in the entire enclosed structure/building from the start of application until application is complete and one of the following ventilation criteria (providing outdoor air) is met: (1) 10 air exchanges; (2) 2 hours of fans or other mechanical ventilation providing outdoor air; (3) 4 hours of vents, windows, or other passive ventilation; (4) 11 hours with no ventilation followed by 1 hour of mechanical ventilation; (5) 11 hours of no ventilation followed by 2 hours of passive ventilation; or (6) 24 hours with no ventilation. After ventilation criteria are met and until the REI expires, do not enter or allow worker entry into treated areas, except as provided in the WPS. Note: after the expiration of the REI whenever etridiazole-treated soil or planting media is being handled or disturbed indoors, continuous ventilation of the area is required at a minimum rate of one complete air exchange per hour.

**EXCEPTION:** If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

## **Chemigation Information**

Chemigation is prohibited in the State of California.

## **Chemigation Instructions**

Pesticide labels contain Directions for Use that are necessary for effecting the purpose for which the product is intended. Further, they protect health and environment

from risks of pesticide contamination of ground water and decrease direct human exposure to pesticide-treated irrigation water.

Apply this product only through pressurized drench (flood), sprinkler, or drip (trickle) irrigation systems. Do not apply this product through any other type of irrigation systems.

Precaution: Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect any irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place.

A person knowledgeable of the chemigation system and responsible for its operation or under the supervision of the responsible person shall shut the system down and make necessary adjustments.

## **Pressurized Drench (Flood) System**

Systems utilizing a pressurized water and pesticide injection system must meet the following requirements:

- Systems using a gravity flow pesticide dispensing system must meter the pesticide into the water at the head of the field and downstream of a hydraulic discontinuity including a drop structure or weir box to decrease potential for water source contamination from backflow if water flow stops.
- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlocked to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, including a positive displacement injection pump (e.g., diaphragm pump), effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- Referring to the rate information contained in this label, apply the appropriate volume of fungicide solution to the crop.
- Upon completing the application of the appropriate volume of Banrot® 40 WP, allow the irrigation system to run long enough to flush the irrigation lines clear of fungicide solution. Do not apply more clear water than is necessary to flush the lines or poor disease control may result.
- Do not apply when wind speed favors drift beyond the area intended for treatment.

### **Drip (Trickle) Chemigation**

- The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
- The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
- The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- Systems must use a metering pump, including a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

### **Product Information**

Pesticide supply tanks are recommended for the application of this product. See label instructions for dilution use rates and timing of application. Agitate continuously during application.

### **Ornamentals and Nursery Crops**

**Greenhouse Restrictions: For use in commercial greenhouses only; use in residential greenhouses or other indoor plant sites is prohibited.**

- Irrigate the crop prior to applying Banrot® 40 WP in order to ensure even moisture content over the entire crop.
- Apply Banrot 40 WP at the appropriate concentration for the crop and disease(s) targeted.

Banrot 40 WP is a soil fungicide for the control of damping-off, root and stem rot diseases caused by Pythium, Phytophthora, Rhizoctonia, Fusarium, and Thielaviopsis. It may be used at time of seeding and transplanting as a drench to control the labeled pathogens in commercial greenhouses and outdoor culture, including shade houses.

Constantly agitate Banrot 40 WP suspensions in water to assure uniform mixing and application. Banrot 40 WP suspensions need not be protected from light. Use at specified rates and concentrations. Overdosage may result in toxicity to sensitive plants. If plants are suspect of being sensitive, apply fungicide to a small number to insure safety. Do not mix Banrot 40 WP with other pesticides or nutrients unless adequately tested first.

Thoroughly rinse Banrot 40 WP residues out of all equipment after use.

**IMPORTANT:** If, after using Banrot® 40 WP as directed, treatment is not effective, a resistant strain of fungus may be present. Consult with your local Everris representative or distributor, your State Agricultural Experiment Station or your State Agricultural Extension Service for advice on the prompt use of some other labeled fungicide.

## Resistance Management

For resistance management, please note that Banrot 40 WP contains both a Group 1 (thiophanate-methyl) and Group 14 (etr Diazole) fungicide. Any fungal population may contain individuals naturally resistant to Banrot 40 WP and other Group 1 or Group 14 fungicides. A gradual or total loss of pest control may occur over time if these fungicides are used repeatedly in the same fields or other use sites. Appropriate resistance-management strategies should be followed.

To delay fungicide resistance, take one or more of the following steps:

- Rotate the use of Banrot 40 WP or other Group 1 or 14 fungicides within a growing season sequence with different groups that control the same pathogens.
- Use tank mixtures with fungicides from a different group that are equally effective on the target pest when such use is permitted. Use at least the minimum application rate as labeled by the manufacturer.
- Adopt an integrated disease management program for fungicide use that includes scouting, use of historical information related to pesticide use, and crop rotation, and which considers host plant resistance, impact of environmental conditions such as disease development, disease thresholds, as well as cultural, biological and other chemical control practices.
- Where possible, make use of predictive disease models to effectively time fungicide applications. Note that using predictive models alone is not sufficient to manage resistance.
- Monitor treated fungal populations for resistance development.
- Contact your local extension specialist or certified crop advisor for any additional pesticide resistance-management and/or IPM recommendations for specific crops and pathogens.
- For further information or to report suspected resistance contact Everris at 1-800-492-8255. You can also contact your pesticide distributor or university extension specialist to report resistance.

PLANT TYPE	APPLICATION INSTRUCTIONS
<p><b>BEDDING PLANTS</b></p> <p>Ageratum Ajuga Alyssum Aster Balsam Begonia Carnation Celosia Coleus Dahlia Dusty Miller Gynura Impatiens Larkspur Marigold Nasturtiums Pansy</p> <p>Petunia Phlox Salvia Snapdragon Verbena Zinnia</p>	<p><b>BEDDING PLANTS</b> (soil 2" to 3" deep). Mix 4-8 oz. of Banrot® 40 WP with 100 gal. of water and apply to 800 sq. ft. of bed area (equivalent to 1 pint/sq. ft. of soil surface) and use at time of seeding and transplanting. Irrigate immediately with additional water equal to at least half the volume of the fungicidal drench for improved soil penetration of the fungicide. Re-treat at 4 to 8 week intervals if additional applications are necessary.</p> <p><b>Use Restrictions:</b></p> <ul style="list-style-type: none"> <li>• Minimum retreatment interval (RTI) is 28 days (4 weeks). Do not make more than 13 applications of product per year.</li> <li>• Do not exceed maximum yearly application rate of 354 lb. of product/acre (53 lb/acre of etridiazole and 88.5 lb/acre of thiophanate-methyl) on bedding sites.</li> </ul>
<p><b>FOLIAGE PLANTS</b></p> <p>Aglaonema Aralia Cactus Caladium Dieffenbachia Dracaena Epipremnum Fern Ficus Fittonia Hoya Maranta Nautilocalyx Peperomia Philodendron Pilea Schefflera Syngonium Tradescantia</p>	<p><b>BENCH PLANTINGS OF TROPICAL FOLIAGE PLANTS</b> (peat and other high organic growth media 6" deep or more). Mix 6-12 oz. of Banrot 40 WP with 100 gal. of water and apply to 400 sq. ft. of bench area (equivalent to 1 quart/sq. ft. of soil surface). Irrigate immediately with additional water equal to at least half the volume of the fungicidal drench for improved soil penetration of the fungicide. Re-treat at 8 to 12 week intervals to maintain disease control.</p> <p><b>Use Restrictions:</b></p> <ul style="list-style-type: none"> <li>• Minimum retreatment interval (RTI) is 56 days (8 weeks). Do not make more than 6 applications of product per year.</li> <li>• Do not exceed maximum yearly application rate of 490 lb. of product/acre (73.5 lb/acre of etridiazole and 122.5 lb/acre of thiophanate-methyl) for tropical foliage bench plantings.</li> </ul> <p><b>BED-GROWN PLANTS</b> Mix 6-12 oz. of Banrot 40 WP with 100 gal. of water per 400 sq. ft. (equivalent to 1 quart/sq. ft. soil surface) or apply in sufficient volume to saturate the soil mixture. Re-treat at 4 to 12 week intervals if necessary.</p>

**PLANT TYPE (continued)****APPLICATION INSTRUCTIONS (continued)****BED-GROWN PLANTS (continued)****Use Restrictions:**

- Minimum retreatment interval (RTI) is 28 days (4 weeks). Do not make more than 13 applications of product per year.
- Do not exceed maximum yearly application rate of 1,062 lb. of product/acre (159 lb/acre of etridiazole and 265.5 lb/acre of thiophanate-methyl) for tropical foliage bed-grown plants.

**CONTAINER & BED-GROWN****CONTAINER & BED-GROWN PLANTS**

Mix 6-12 oz. of Banrot® 40 WP with 100 gal. of water per 400 sq. ft. (equivalent to 1 quart/sq. ft. soil surface) or apply in sufficient volume to saturate the soil mixture, including ½ pint per 6" pot. Irrigate immediately with additional water equal to at least half the volume of the fungicidal drench for improved soil penetration of the fungicide. Re-treat at 4 to 12 week intervals if necessary with a 6-12 oz. per 100 gal. rate.

**Use Restrictions:**

- Minimum retreatment interval (RTI) is 28 days (4 weeks). Do not make more than 13 applications of product per year.
- Do not exceed maximum yearly application rate of 1,062 lb. of product/acre (159 lb/acre of etridiazole and 265.5 lb/acre of thiophanate-methyl) for container and bed-grown herbaceous and woody plants.

**HERBACEOUS**

African Violet	Gypsophila
Begonia	Ivy
Calendula	Kalanchoe
Carnation	Lobelia
Chrysanthemum	Poinsettia
Coleus	Statice
Dahlia	Zygocactus

**WOODY**

Acer	Ligustrum
Aucuba	Photinia
Azalea	Pine
Boxwood	Pittosporum
Camellia	Podocarpus
Cedar	Rhododendron
Chamaedorea Palm	Spruce
Crape Myrtle	Taxus
Croton	Thuja

Euonymus  
Fatsia  
Ficus  
Fir  
Gardenia  
Hydrangea  
Ilex  
Ixora  
Juniper  
Leucothe

**PLANT TYPES (continued)****APPLICATION INSTRUCTIONS (continued)****COFFEE SEEDLINGS (Propagation)****SEED BEDS**

**Prior to sowing seed:** Mix 4 ounces of Banrot® 40 WP with 50 gal. of water and apply 1 gal. per 1.2 sq. yd. two days prior to sowing seeds, thoroughly wetting the planting medium.

**Use Restrictions:**

- Apply once prior to sowing seed.

**Seeds previously sown:** Mix 3 oz. of Banrot 40 WP with 50 gal. of water and apply 1 gal. per 1.2 sq. yd. If seedlings have emerged, follow application immediately by irrigating with additional water equal to at least ½ the volume of the fungicidal drench for improved soil penetration and to rinse foliage. Re-treat at 4 - 12 week intervals if necessary.

**Use Restrictions:**

- For non-bearing seedlings only. Do not apply after field transplanting and within two years of the first coffee bean harvest.
- Do not apply to bearing coffee plants. Preharvest interval (PHI) for coffee is 2 years (730 days).
- Minimum retreatment interval (RTI) is 28 days (4 weeks). Do not exceed 4 applications of product per year.
- Do not exceed maximum yearly application rate of 65.5 lb. of product/acre (9.9 lb/acre of etridiazole and 17 lb./acre of thiophanate-methyl) for coffee seeds previously sown.

**CONTAINER SEEDLINGS**

Mix 6 ounces of Banrot 40 WP with 50 gal. of water. Apply 1.7 fl. oz. of this mixture to a 1 qt. container. Follow-up application by immediately irrigating with additional water equal to at least ½ the volume of the fungicidal drench for improved soil penetration and to rinse foliage. Re-treat at 4 to 12 week intervals if necessary.

**Use Restrictions:**

- For non-bearing seedlings only. Do not apply after field transplanting and within two years of the first coffee bean harvest.
- Do not apply to bearing coffee plants. Preharvest interval (PHI) for coffee is 2 years (730 days).
- Minimum retreatment interval (RTI) is 28 days (4 weeks). Do not exceed 13 applications of product per year.
- Do not exceed maximum yearly application rate of 508 lb. of product/acre (76.2 lb/acre of etridiazole and 127 lb/acre of thiophanate-methyl) for container grown coffee seedlings.



**PLANT TYPES (continued)****APPLICATION INSTRUCTIONS (continued)****CONTAINERIZED CITRUS SEEDLINGS AND LINERS (Propagation)****CONTAINERIZED SEEDLINGS AND LINERS**

**Bench area:** Mix 6-10 oz. of Banrot® 40 WP in 100 gal. water and apply to 800 sq. ft. of bench area (1 pint of solution per sq. ft.) at 4 weeks and 8 weeks after seeding. With seedlings scheduled for planting in ground liner beds out-of-doors, apply an additional drench, in the greenhouse, 1 to 2 weeks prior to field liner bed planting.

**Transplanting:** With plants scheduled for replanting to larger containers in greenhouse culture for the budding process, apply an additional drench upon transplanting. Mix 6-10 oz. Banrot 40 WP with 100 gal. of water per 400 sq. ft. Re-treat at 6 to 12 week intervals if necessary. Follow all applications by immediately irrigating with additional water equal to at least ½ the volume of the fungicidal drench for improved soil penetration and to rinse foliage.

**Use Restrictions:**

- Discontinue any applications 2 months prior to grove planting.
- For non-bearing seedlings and liner use only. Do not apply after field transplanting and within two years of first citrus fruit harvest.
- Do not apply to fruit bearing citrus trees. Preharvest interval (PHI) for citrus is 2 years (730 days).
- Minimum retreatment interval (RTI) is 42 days (6 weeks). Do not apply more than 8 applications of product per year.
- Do not exceed maximum yearly application rate of 545 lb. of product/acre (82 lb/acre of etridiazole and 137 lb/acre of thiophanate-methyl) for containerized citrus seedling and liner transplants.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage and disposal.

**Storage:** Not sensitive to light. Stable at normal temperatures and humidities. Store in a secure dry place at temperature above 45 ° F (7 ° C) in a well-closed container.

**Pesticide Disposal:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**Container Handling:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling, if available. Dispose of empty container in a sanitary landfill or by incineration or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

### DISCLAIMER AND LIMITATION OF LIABILITY

IMPORTANT NOTICE FROM EVERRIS NA INC. ("Everris")  
PLEASE READ BEFORE USE.

To the extent consistent with applicable law, user or buyer accepts the conditions, disclaimer of warranties and limitations of liability. Read the entire directions for use, conditions of warranties and limitations of liability before using this product. If terms are not acceptable, return the unopened product container at once for full refund.

**CONDITIONS:** The directions for use of this product are believed to be adequate and the user or buyer must always follow the label directions carefully and exercise judgment and caution when using this product under their growing conditions. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness, unsatisfactory or substandard results or other unintended consequences may result because of such factors as weather conditions, presence or absence of other materials, or the manner of use or application, all of which are beyond the control of Everris. All such risks shall be assumed by the user or buyer.

**WARRANTY:** This product corresponds to all claims and descriptions set forth on the label and is reasonably fit for the purposes set forth in the directions for use on the label when used in accordance with those directions. This warranty is subject to the conditions and limitations stated herein. Everris recognizes that the rights and remedies of the user or buyer are subject to the provisions of the applicable law, but makes no other warranties or representations, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of

Everris is authorized to make any warranties beyond those contained herein or to modify the warranties contained therein. Subject to the user's or buyer's rights and remedies under the applicable law, Everris disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

**LIMITATIONS OF LIABILITY:** Subject to the user's or buyer's rights and remedies under the applicable law, the exclusive remedy of the user or buyer and the liability of Everris or its affiliates, for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid by the user or Buyer for the quantity of this product involved or, at Everris' election, the replacement of the product. To the extent consistent with applicable law, Everris must have prompt notice of any claim so that a timely investigation of buyer's or user's claim can be made. Buyer and all users shall promptly notify Everris of any claims, whether based on contract, negligence, strict liability, other tort or otherwise or be barred from any remedy.

Manufactured for:  
Everris NA Inc.  
PO Box 3310  
Dublin, OH 43016  
1-800-492-8255

